
GUIDE TO DIARRHEA IN PIGS HOW TO MONITOR AND FIND SOLUTIONS











Diarrhea scoring guide

	Normal	Moderate	Severe
	<2	2-2.5	>2.5
% diarrhea	<10%	10-20%	>20%
	<3 days	3-5 days	>5 days
Actions	Support therapy Acciffiers, eletrolytes Monitor affected animals	Isolate individuals Monitor daily Gruel feed Medicate with injectables Seek veterinarian advice for individual medication Support therapy	Check environmental factors Check feeds and feeding Check disease status Medicate, if needed Seek veterinarian advice for individual medication Support therapy

Fecal score











How to monitor and score

- Fecal score is rated in early morning by looking at the fecal consistency in each pen.
 The fecal scoring guide (1 to 4) is used to gauge the general consistency.
- Typically, assessments by 2-3 people can be averaged to represent the pen fecal score.

 Should be done daily until the fecal score normalizes.

= <u>Total number of pigs with fecal score "4" x 100</u> Total pigs in the pen

Duration of diarrhea

Total days from weaning where pigs exhibit fecal score "4"

Environmental and management causes

- Poor hygiene and sanitation
- Too cold and drafty
- Wetpens
- AIAO failure
- High temperature variation during 24-hour period
- Mixing animals of different ages and/or different sources
- Low colostrum intake
- High percentage of first parity sows in the herd
- Low birth and weaning weights









Nutritional causes

- High crude protein in the
- High fats and oils
- Mycotoxins
- Inappropriate feed ingredients
- Delayed feed/late eaters
- High acid-binding capacity in the diet
- Feed anti-nutritional factors
 High coliforms in drinking water

FECAL CONSISTENCY



